

# Nested Quantifiers

Section 1.5


# Section Summary

- Nested Quantifiers
- Order of Quantifiers
- Translating to and from English
- Negating Nested Quantifiers

# Nested Quantifiers

- Two qualifiers are **nested** if one is within the scope of the other.

**Example:** “Every real number has an inverse” is



$\forall x Q(x)$

$Q(x)$  is  $\exists y P(x,y)$

$P(x,y)$  is “ $x+y=0$ ”

where the domains of  $x$  and  $y$  are the real numbers.

# Thinking of Nested Quantification as Nested Loops

Loop through all values of  $x$ . At each step, loop through all values of  $y$ .

- $\forall x \forall y P(x,y)$ 
  - If  $P(x,y)$  is false for some pair of  $x$  and  $y$ , then  $\forall x \forall y P(x,y)$  is false and both the outer and inner loop terminate.
  - $\forall x \forall y P(x,y)$  is true if the outer loop ends after stepping through each  $x$ .
- $\forall x \exists y P(x,y)$ 
  - The inner loop ends when a pair  $x$  and  $y$  is found such that  $P(x,y)$  is true.
  - If no  $y$  is found such that  $P(x,y)$  is true, the outer loop terminates as  $\forall x \exists y P(x,y)$  has been shown to be false.
  - $\forall x \exists y P(x,y)$  is true if the outer loop ends after stepping through each  $x$ .

# Order of Quantifiers

## Examples:

1.  $P(x,y) : "x + y = y + x."$  Assume that  $U$  is the real numbers.
  - $\forall x \forall y P(x,y)$  and  $\forall y \forall x P(x,y)$  have the same truth value.
  - $\exists x \exists y P(x,y)$  and  $\exists y \exists x P(x,y)$  have the same truth value.
2.  $Q(x,y) : "x + y = 0."$  Assume that  $U$  is the real numbers.
  - $\forall x \exists y Q(x,y)$  is true, but
  - $\exists y \forall x Q(x,y)$  is false

# Questions on Order of Quantifiers

**Example 1:** Let  $U$  be the real numbers,

Define  $P(x,y)$ : “ $x \cdot y = 0$ ”

What is the truth value of the following:

1.  $\forall x \forall y P(x,y)$

Answer: **False**

2.  $\forall x \exists y P(x,y)$

Answer: **True**

3.  $\exists x \forall y P(x,y)$

Answer: **True**

4.  $\exists x \exists y P(x,y)$

Answer: **True**

# Questions on Order of Quantifiers

**Example 2:** Let  $U$  be the real numbers,

Define  $P(x,y) : "x / y = 1"$

What is the truth value of the following:

1.  $\forall x \forall y P(x,y)$

Answer: **False**

2.  $\forall x \exists y P(x,y)$

Answer: **True**

3.  $\exists x \forall y P(x,y)$

Answer: **False**

4.  $\exists x \exists y P(x,y)$

Answer: **True**

# Quantifications of Two Variables

Statement	When True?	When False
$\forall x \forall y P(x, y)$ $\forall y \forall x P(x, y)$	$P(x,y)$ is true for every pair $x,y$ .	There is a pair $x, y$ for which $P(x,y)$ is false.
$\forall x \exists y P(x, y)$	For every $x$ there is a $y$ for which $P(x,y)$ is true.	There is an $x$ such that $P(x,y)$ is false for every $y$ .
$\exists x \forall y P(x, y)$	There is an $x$ for which $P(x,y)$ is true for every $y$ .	For every $x$ there is a $y$ for which $P(x,y)$ is false.
$\exists x \exists y P(x, y)$ $\exists y \exists x P(x, y)$	There is a pair $x, y$ for which $P(x,y)$ is true.	$P(x,y)$ is false for every pair $x,y$



# Translating Nested Quantifiers into English

Let  $U$  be all students in your school. Using  $C(x)$  = “ $x$  has a computer,” and  $F(x,y)$  = “ $x$  and  $y$  are friends,” translate the following statements.

- $\forall x (C(x) \vee \exists y (C(y) \wedge F(x, y)))$

**Solution:** Every student in your school has a computer or has a friend who has a computer.

- $\exists x \forall y \forall z ((F(x, y) \wedge F(x, z) \wedge (y \neq z)) \rightarrow \neg F(y, z))$

**Solution:** There is a student none of whose friends are also friends with each other.

# Translating Mathematical Statements into Predicate Logic

1. Rewrite the statement to **make the implied quantifiers and domains explicit**
2. Introduce **variables** and specify the **domain** for them
3. Rewrite the statement using quantifiers, variables, and logic expressions.

**Example :** Translate “The sum of two positive integers is always positive” into a logical expression.

1. “For every two positive integers, the sum of these integers is positive.”
2. “For all positive integers  $x$  and  $y$ ,  $x + y$  is positive.”
3.  $\forall x \forall y (x + y > 0)$ , where the domain of both variables consists of all positive integers

# Translating English into Logical Expressions Example

**Example:** Use quantifiers to express the statement “There is a **woman** who has taken a **flight** on every **airline**.”

**Solution:**

- Let  $P(w,f)$  = “ $w$  has taken  $f$ ” and  $Q(f,a)$  = “ $f$  is a flight on  $a$ .”
- The domain of  $w$  is all women, the domain of  $f$  is all flights, and the domain of  $a$  is all airlines.
- Then the statement can be expressed as:

$$\exists w \forall a \exists f (P(w,f) \wedge Q(f,a))$$

# Questions on Translation from English

Choose the obvious predicates and express in predicate logic.

**Example 1:** “Brothers are siblings.”

**Solution:**  $\forall x \forall y (B(x,y) \rightarrow S(x,y))$

**Example 2:** “Siblinghood is symmetric.”

**Solution:**  $\forall x \forall y (S(x,y) \rightarrow S(y,x))$

**Example 3:** “Everybody loves somebody.”

**Solution:**  $\forall x \exists y L(x,y)$

**Example 4:** “There is someone who is loved by everyone.”

**Solution:**  $\exists y \forall x L(x,y)$

**Example 5:** “There is someone who loves someone.”

**Solution:**  $\exists x \exists y L(x,y)$

**Example 6:** “Everyone loves himself”

**Solution:**  $\forall x L(x,x)$

# Negating Nested Quantifiers

**Example 1:** Express the negation of the statement  $\forall x \exists y (xy=1)$  so that no negation precedes a quantifier.

**Solution:** Use **De Morgan's Laws** to move the negation as far inwards as possible.

1.  $\neg \forall x \exists y (xy = 1)$
2.  $\exists x \neg \exists y (xy = 1)$  by De Morgan's for  $\forall$
3.  $\exists x \forall y \neg (xy = 1)$  by De Morgan's for  $\exists$
4.  $\exists x \forall y (xy \neq 1)$

# Negating Nested Quantifiers

Translate the following statement into a logical expression.

**“There does not exist a woman who has taken a flight on every airline.”**

**Solution:**

- Translate the positive sentence into a logical expression
  - $\exists w \forall a \exists f (P(w,f) \wedge Q(f,a))$  [by previous example]
  - $P(w,f)$ : “ $w$  has taken  $f$ ”     $Q(f,a)$ : “ $f$  is a flight on  $a$ .”
- Find the negation of the logical expression
  - $\neg \exists w \forall a \exists f (P(w,f) \wedge Q(f,a))$
  - $\forall w \neg \forall a \exists f (P(w,f) \wedge Q(f,a))$
  - $\forall w \exists a \neg \exists f (P(w,f) \wedge Q(f,a))$
  - $\forall w \exists a \forall f \neg (P(w,f) \wedge Q(f,a))$
  - $\forall w \exists a \forall f (\neg P(w,f) \vee \neg Q(f,a))$

“For every woman there is an airline such that for all flights, this woman has not taken that flight or that flight is not on this airline”