

Binomial Coefficients and Identities

Section 6.4

Section Summary

- The Binomial Theorem
- Pascal's Identity and Triangle

Powers of Binomial Expressions

Definition: A *binomial* expression is the sum of two terms, such as $x + y$. (More generally, these terms can be products of constants and variables.)

- We can use counting principles to find the **coefficients** in the expansion of $(x + y)^n$ where n is a positive integer.

Example: Expand $(x + y)^3$
 $x^3 + 3x^2y + 3xy^2 + y^3$

continued →

Powers of Binomial Expressions

$$\begin{aligned}(x+y)^3 &= (x+y)(x+y)(x+y) \\ &= (xx + xy + yx + yy)(x+y) \\ &= xxx + xxy + xyx + xyy + yxx + yxy + yyx + yyy \\ &= x^3 + 3x^2y + 3xy^2 + y^3\end{aligned}$$

How many ways are there to get the terms

| | | |
|--------|-------------------------------|-----------------------------|
| x^3 | : Choose x three times | $\binom{3}{3} = C(3,3) = 1$ |
| x^2y | : Choose x twice (and y once) | $\binom{3}{2} = C(3,2) = 3$ |
| xy^2 | : Choose x once (and y twice) | $\binom{3}{1} = C(3,1) = 3$ |
| y^3 | : Choose x zero times | $\binom{3}{0} = C(3,0) = 1$ |

Binomial Theorem

Binomial Theorem: Let x and y be variables, and n a nonnegative integer. Then:

$$(x+y)^n = \sum_{j=0}^n \binom{n}{j} x^{n-j} y^j = \binom{n}{0} x^n + \binom{n}{1} x^{n-1} y + \cdots + \binom{n}{n-1} x y^{n-1} + \binom{n}{n} y^n.$$

Proof: We use combinatorial reasoning. The terms in the expansion of $(x + y)^n$ are of the form $x^{n-j}y^j$ for $j = 0, 1, 2, \dots, n$. To form the term $x^{n-j}y^j$, it is necessary to choose $n-j$ x 's from the n sums. Therefore, the coefficient of $x^{n-j}y^j$ is $\binom{n}{n-j}$, which equals $\binom{n}{j}$. ◀

Using the Binomial Theorem

Example: What is the expansion of $(x + y)^4$?

Solution: From the binomial theorem, we know that

$$\begin{aligned}(x + y)^4 &= \sum_{j=0}^4 \binom{4}{j} x^{4-j} y^j \\ &= \binom{4}{0} x^4 + \binom{4}{1} x^3 y + \binom{4}{2} x^2 y^2 + \binom{4}{3} x y^3 + \binom{4}{4} y^4 \\ &= x^4 + 4x^3 y + 6x^2 y^2 + 4x y^3 + y^4.\end{aligned}$$

Using the Binomial Theorem

Example: What is the coefficient of $x^{12}y^{13}$ in the expansion of $(2x - 3y)^{25}$?

Solution: We view the expression as $(2x + (-3y))^{25}$.
By the binomial theorem

$$(2x + (-3y))^{25} = \sum_{j=0}^{25} \binom{25}{j} (2x)^{25-j} (-3y)^j.$$

The coefficient of $x^{12}y^{13}$ in the expansion is obtained when $j = 13$.

$$\binom{25}{13} 2^{12} (-3)^{13} = -\frac{25!}{13!12!} 2^{12} 3^{13}.$$

A Useful Identity

Corollary 1: With $n \geq 0$, $\sum_{k=0}^n \binom{n}{k} = 2^n$.

Proof (using *binomial theorem*): With $x = 1$ and $y = 1$, from the binomial theorem we see that:

$$2^n = (1 + 1)^n = \sum_{k=0}^n \binom{n}{k} 1^k 1^{(n-k)} = \sum_{k=0}^n \binom{n}{k}.$$




A Useful Identity

Corollary 1: With $n \geq 0$, $\sum_{k=0}^n \binom{n}{k} = 2^n$.

Proof (combinatorial): Consider the subsets of a set with n elements. There are $\binom{n}{0}$ subsets with zero elements, $\binom{n}{1}$ with one element, $\binom{n}{2}$ with two elements, ..., and $\binom{n}{n}$ with n elements. Therefore the total is $\sum_{k=0}^n \binom{n}{k}$.

Since, we know that a set with n elements has 2^n subsets, we conclude: $\sum_{k=0}^n \binom{n}{k} = 2^n$.



Pascal's Identity

Pascal's Identity: If n and k are integers with $n \geq k \geq 0$, then

$$\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}.$$

Proof (combinatorial): Let T be a set where $|T| = n + 1$, $a \in T$, and $S = T - \{a\}$. There are $\binom{n+1}{k}$ subsets of T containing k elements. Each of these subsets either:

- contains a with $k - 1$ other elements, or
- contains k elements of S and not a .

There are


- $\binom{n}{k-1}$ subsets of k elements that contain a , since there are $\binom{n}{k-1}$ subsets of $k - 1$ elements of S ,
- $\binom{n}{k}$ subsets of k elements of T that do not contain a , because there are $\binom{n}{k}$ subsets of k elements of S .

Hence, $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$. ◀

*See Exercise 19
for an algebraic
proof.*


Pascal's Triangle

The n th row in the triangle consists of the binomial coefficients $\binom{n}{k}$, $k = 0, 1, \dots, n$.



By Pascal's identity:

$$\binom{6}{4} + \binom{6}{5} = \binom{7}{5}$$



By Pascal's identity, adding two adjacent binomial coefficients results in the binomial coefficient in the next row between these two coefficients.

Pascal's Triangle

$$\binom{0}{0}$$

$$\binom{1}{0} \binom{1}{1}$$

$$\binom{2}{0} \binom{2}{1} \binom{2}{2}$$

$$\binom{3}{0} \binom{3}{1} \binom{3}{2} \binom{3}{3}$$

$$\binom{4}{0} \binom{4}{1} \binom{4}{2} \binom{4}{3} \binom{4}{4}$$

$$\binom{5}{0} \binom{5}{1} \binom{5}{2} \binom{5}{3} \binom{5}{4} \binom{5}{5}$$

$$\binom{6}{0} \binom{6}{1} \binom{6}{2} \binom{6}{3} \binom{6}{4} \binom{6}{5} \binom{6}{6}$$

$$\binom{7}{0} \binom{7}{1} \binom{7}{2} \binom{7}{3} \binom{7}{4} \binom{7}{5} \binom{7}{6} \binom{7}{7}$$

$$\binom{8}{0} \binom{8}{1} \binom{8}{2} \binom{8}{3} \binom{8}{4} \binom{8}{5} \binom{8}{6} \binom{8}{7} \binom{8}{8}$$

By Pascal's identity:

$$\binom{6}{4} + \binom{6}{5} = \binom{7}{5}$$

1

1 1

1 2 1

1 3 3 1

1 4 6 4 1

1 5 10 10 5 1

1 6 15 20 15 6 1

1 7 21 35 35 21 7 1

1 8 28 56 70 56 28 8 1

$$(x+y)^6 = 1x^6 + 6x^5y + 15x^4y^2 + 20x^3y^3 + 15x^2y^4 + 6xy^5 + 1y^6$$

Pascal's Triangle

$$\binom{0}{0}$$

$$\binom{1}{0} \binom{1}{1}$$

$$\binom{2}{0} \binom{2}{1} \binom{2}{2}$$

$$\binom{3}{0} \binom{3}{1} \binom{3}{2} \binom{3}{3}$$

$$\binom{4}{0} \binom{4}{1} \binom{4}{2} \binom{4}{3} \binom{4}{4}$$

$$\binom{5}{0} \binom{5}{1} \binom{5}{2} \binom{5}{3} \binom{5}{4} \binom{5}{5}$$

$$\binom{6}{0} \binom{6}{1} \binom{6}{2} \binom{6}{3} \binom{6}{4} \binom{6}{5} \binom{6}{6}$$

$$\binom{7}{0} \binom{7}{1} \binom{7}{2} \binom{7}{3} \binom{7}{4} \binom{7}{5} \binom{7}{6} \binom{7}{7}$$

$$\binom{8}{0} \binom{8}{1} \binom{8}{2} \binom{8}{3} \binom{8}{4} \binom{8}{5} \binom{8}{6} \binom{8}{7} \binom{8}{8}$$

By Pascal's identity:

$$\binom{6}{4} + \binom{6}{5} = \binom{7}{5}$$

1

1 1

1 2 1

1 3 3 1

1 4 6 4 1

1 5 10 10 5 1

1 6 15 20 15 6 1

1 7 21 35 35 21 7 1

1 8 28 56 70 56 28 8 1

$$(x+y)^6 = 1x^6 + 6x^5y + 15x^4y^2 + 20x^3y^3 + 15x^2y^4 + 6xy^5 + 1y^6$$